Anti- PAK1/2/3 (Phospho-Thr423/402/421)Polyclonal



 Catalog No.
 Size

 A100074-01
 50 μl

 A100074-02
 100 μl



Specificity Anti- PAK1/2/3 (Phospho-Thr423/402/421) (human mouse rat)

Source Rabbit Polyclonal

Application ELISA IHC

Form Liquid, 1 mg/ml

Pruduct

Swiss-Prot No.: Q13153/Q13177/O75914

Other Names: EC 2.7.11.1, Gamma-PAK, P21-activated kinase 2, PAK 2, PAK-2, PAK65, PAKI, S6/H4 kinase, kinase PAK2, p21-activated kinase 2, p21-activated protein kinase IBeta-PAK, CDC42/RAC effector kinase PAK-B, EC 2.7.1., EC 2.7.11.1, OPHN3, Oligophrenin-3, P65-PAK, PAK 3, PAK-3, PAKB, STK4, Serine/threonine-protein kinase PAK 3, kinase PAK3, p21-activated kinase 3

Specificity and Sensitivity

PAK1/2/3 (Phospho-Thr423/402/421) antibody detects endogenous levels of PAK1/2/3 only when phosphorylated at threonine 423/402/421.

Source and Purification

The antiserum was produced against synthesized phosphopeptide derived from human PAK1/2/3 around the phosphorylation site of threonine 423/402/421 (R-S-T^P-M-V). The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

Application Notes

Optimal working dilutions should be determined experimentally by the investigator. Suggested starting dilutions are as follows:

IHC: 1:50~1:100 ELISA: 1:5000


Storage Buffer

Rabbit IgG in phosphate buffered saline (without Mg^{2+} and Ca^{2+}), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.

Storage Instructions

Stable for 1 year at -20°C from date of shipment. For maximum recovery of product, centrifuge the original vial after thawing and prior to removing the cap. Aliquot to avoid repeated freezing and thawing. Aliquot will be stable at 4°C for 3 months.

Images



P-peptide -

Immunohistochemistry analysis of paraffin-embedded human breast carcinoma tissue using PAK1/2/3 (Phospho- Thr423/ 402/421) antibody.

