

Anti- PKC θ (Phospho-Ser695) Polyclonal Antibody



| <u>Catalog No.</u> | <u>Size</u> |
|--------------------|-------------|
| A100104-01 | 50 μ l |
| A100104-02 | 100 μ l |

| | |
|--------------------|--|
| Specificity | Anti- PKC θ (Phospho-Ser695) (human mouse) |
| Source | Rabbit Polyclonal |
| Application | WB ELISA IHC |
| Form | Liquid, 1 mg/ml |

Product

Swiss-Prot No.: Q04759

Other Names: EC 2.7.11.13; kinase PKC-theta; KPCT; nPKC-theta; PKC-theta; PKCQ; PRKCQ; PRKCT; Protein kinase C, theta type

Specificity and Sensitivity

PKC θ (Phospho-Ser695) antibody detects endogenous levels of PKC θ only when phosphorylated at serine 695.

Source and Purification

The antiserum was produced against synthesized phosphopeptide derived from human PKC θ around the phosphorylation site of serine 695 (N-F-S^P-F-M).

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

Application Notes

Optimal working dilutions should be determined experimentally by the investigator. Suggested starting dilutions are as follows:

WB: 1:500~1:3000 IHC: 1:50~1:100 ELISA: 1:5000


Storage Buffer

Rabbit IgG in phosphate buffered saline (without Mg²⁺ and Ca²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.


Storage Instructions

Stable for 1 year at -20°C from date of shipment. For maximum recovery of product, centrifuge the original vial after thawing and prior to removing the cap. Aliquot to avoid repeated freezing and thawing. Aliquot will be stable at 4°C for 3 months.

Images



P-peptide - +
Immunohistochemistry analysis of paraffin-embedded human breast carcinoma tissue using PKC θ (Phospho-Ser695) antibody.



Western blot analysis of extracts from Jurkat cells, treated with EGF (200ng/ml, 15mins), using PKC θ (Phospho-Ser695) antibody.