

# Anti- Src (Phospho-Tyr529) Polyclonal Antibody



<u>Catalog No.</u>	<u>Size</u>
A100190-01	50 µl
A100190-02	100 µl

<b>Specificity</b>	Anti- Src (Phospho-Tyr529) (human mouse rat)
<b>Source</b>	Rabbit Polyclonal
<b>Application</b>	WB ELISA IHC
<b>Form</b>	Liquid, 1 mg/ml

## Product

**Swiss-Prot No.:** P12931

## Specificity and Sensitivity

Src (phospho-Tyr529) antibody detects endogenous levels of Src only when phosphorylated at tyrosine 529.

## Source and Purification

The antiserum was produced against synthesized phosphopeptide derived from human Src around the phosphorylation site of tyrosine 529 (P-Q-YP-Q-P).

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

## Application Notes

Optimal working dilutions should be determined experimentally by the investigator. Suggested starting dilutions are as follows:

WB: 1:500~1:3000    IHC: 1:50~1:100    ELISA: 1:10000


## Storage Buffer

Rabbit IgG in phosphate buffered saline (without Mg<sup>2+</sup> and Ca<sup>2+</sup>), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.

## Storage Instructions


Stable for 1 year at -20°C from date of shipment. For maximum recovery of product, centrifuge the original vial after thawing and prior to removing the cap. Aliquot to avoid repeated freezing and thawing. Aliquot will be stable at 4°C for 3 months.

## Images



P-peptide                    -                    +

Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue, using Src (phospho- Tyr529) antibody.



P-peptide                    -                    +

Western blot analysis of extracts from COS7 cells, using Src (phospho-Tyr529) antibody.