# Anti-NMDAR2B (Phospho-Tyr1336) Polyclonal Antibody

Catalog No. Size A100310-01 **50** µl A100310-02 100 µl



**Specificity** Anti- NMDAR2B (Phospho-Tyr1336) (human mouse rat)

Source Rabbit Polyclonal WB ELISA IHC **Application** 

Liquid, 1 mg/ml

#### **Product**

**Form** 

Swiss-Prot No.: Q13224

Other Names: Glutamate [NMDA] receptor subunit epsilon 2 precursor; GRIN2B; hNR3; N-methyl D-aspartate receptor subtype 2B; N-methyl-D-aspartate receptor subtype 2B; N-methyl-D-aspartate receptor subunit 3; NMDE2; NME2; NR2B; NR3

# **Specificity and Sensitivity**

NMDAR2B (Phospho-Tyr1336) antibody detects endogenous levels of NMDAR2B only when phosphorylated at tyrosine 1336.

#### **Source and Purification**

The antiserum was produced against synthesized phosphopeptide derived from human NMDAR2B around the phosphorylation site of tyrosine 1336 (S-P-YP-A-H). The antibody was affinity-purified from rabbit antiserum by

affinity-chromatography using epitope-specific immunogen.

## **Application Notes**

Optimal working dilutions should be determined experimentally by the investigator. Suggested starting dilutions are as follows:

WB: 1:500~1:3000 IHC: 1:50~1:100 ELISA: 1:10000


### Storage Buffer

Rabbit IgG in phosphate buffered saline (without Mg<sup>2+</sup> and Ca<sup>2+</sup>), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.

### **Storage Instructions**


Stable for 1 year at -20°C from date of shipment. For maximum recovery of product, centrifuge the original vial after thawing and prior to removing the cap. Aliquot to avoid repeated freezing and thawing. Aliquot will be stable at 4°C for 3 months.

### **Images**



P-peptide

Immunohistochemistry analysis of paraffin-embedded human brain tissue using NMDAR2B (Phospho-Tyr1336) antibody.



Western blot analysis of extracts from Jurkat cells, treated with TNF (20ng/ml, 30mins), using NMDAR2B (Phospho-Tyr1336) antibody.

